

Evaluation of bone mineral density in chronic glue sniffers

M. Ruşen Dünderöz¹, S. Ümit Sarıcı¹, Tümer Türkbay², Barboros Baykal³, Murat Kocaoğlu⁴
H. İbrahim Aydın¹, Erdal Gökçay¹

Departments of ¹Pediatrics, ²Child Psychiatry, ³Orthopedics, and ⁴Radiology, Gülhane Military Medical Academy, Ankara, Turkey

SUMMARY: Dünderöz MR, Sarıcı SÜ, Türkbay T, Baykal B, Kocaoğlu M, Aydın Hİ, Gökçay E. Evaluation of bone mineral density in chronic glue sniffers. Turk J Pediatr 2002; 44: 326-329.

Although acute and chronic toxic effects of inhalant (glue) abuse have been well demonstrated on many organ systems, the effects on the skeletal system and bone mineral content of young people with this addiction have, to our knowledge, not yet been investigated by bone mineral density measurement. In the present study bone mineral density was measured by the dual-energy X-ray absorptiometry method in 25 children and adolescents with inhalant abuse and compared with that of a control group (n=30) to detect whether there was any delay in bone development or any decrease in bone mass. Chronological age, height and weight, serum calcium, phosphorus and alkaline phosphatase levels of the study group were not significantly different from those of the control group ($p>0.05$), whereas bone mineral density was significantly reduced in the study group ($p=0.001$). Teenagers with glue vapor abuse may carry an increased risk of future fracture even though the exact mechanism(s) responsible for the toxicity of glue vapor on bone metabolism remains to be determined. To ascertain the exact component of glue responsible for bone demineralisation may be of value in proposing a change in the composition of the glue. Education and/or rehabilitation programs currently have the greatest importance in preventing and overcoming the harmful effects of this public health problem which is so common in young children and adolescents.

Key words: addiction, adolescent, bone mineral density, children, glue, inhalant abuse.

Inhalant abuse is one of the most common problems of young children and adolescents in developing countries. Desired effects with intentional inhalation of a volatile substance appear to be euphoria, tranquillity, relaxation and hallucination. Glue/adhesive sniffing is the most common form of volatile substance abuses since these substances are different from other drugs in that they are not sold illegally¹.

There are significant morbidity and mortality, both organic and psychosocial, associated with inhalant abuse. Both acute and chronic toxic effects in neuropsychiatric, urological, hematological, cardiovascular, pulmonary and gastrointestinal systems, including death due to cardiac arrhythmia or pulmonary and cerebral edema, encephalopathy syndromes, cerebral damage, optic atrophy, peripheral neuropathy, muscle weakness, gastrointestinal disturbances, or hepatic and renal damage well have been

documented²⁻⁴. However, toxic effects of inhalant abuse on the skeletal system and bone mineral content of young people with this addiction have, to our knowledge, not yet been evaluated by bone mineral density (BMD) measurement although childhood and adolescence are critical periods of bone mineralization and skeletal development.

In the present study, BMD was measured by the dual-energy X-ray absorptiometry (DXA) method in children and adolescents with inhalant abuse and compared with that of a control group to detect whether there was any delay in bone development or any decrease in bone mass.

Material and Methods

This study was performed between September 2000-May 2001 at the Government's Education and Rehabilitation Center of Children in

Ankara, where inpatient and outpatient chemical dependency treatment and prevention programs are conducted. The study group consisted of 25 boys whose ages were between 13 to 19 years (mean 14.9 ± 2.5 years). These cases had overcome the habit at least one month previously (range 1 to 3 months, mean 2.3 months), and the duration of abuse before rehabilitation was between 2 to 5 years (mean 3.2 years). All the cases were smokers, and none took any antioxidant drugs (vitamin C, vitamin E, selenium, etc.) prior to or during the study. The parents and authorities were informed about the study and informed consent was obtained from the cases and/or their parents.

Thirty healthy volunteer adolescents of similar age (mean 15.1 ± 2.9 years) without any addiction were chosen as the control for each case in the study and control groups, brief history was taken, and a complete physical examination was performed. Blood samples for complete blood count, serum calcium, phosphorus and alkaline phosphatase (ALP), and renal and liver function tests were obtained, and a urinalysis was performed. Cases with any signs or symptoms of any acute or chronic illness were excluded from the study.

Body weight and height measurements of the cases in the study were taken carefully by the same experienced personnel. BMD was measured by DXA method^{5,6} at L2-L4 levels of lumbar vertebrae using the DXA Norland (Fort Atkinson, WI, USA) XR-36 densitometer. Spinal BMD was measured in supine position; the scan time ranged between 3 to 5 minutes, and results for spinal BMD measurements were expressed in g/cm^2 .

Mann-Whitney U test was used in statistical analysis of the data.

Results

Chronological age, anthropometric measurements, serum calcium, phosphorus and ALP levels, and BMD measurements of the study and control groups are shown in Table I. Chronological age, height and weight, serum calcium, phosphorus and ALP levels of the study group were not significantly different from those of the control group ($p > 0.05$), whereas BMD was significantly reduced in the study group ($p = 0.001$) (Table I).

Discussion

The effects and toxicity of various components of volatile substances on skeletal development and mineralization have been investigated in only a few studies, most of which have been performed on animals. Subchronic toluene exposure caused inhibition of skeletal growth (torso length, rump width) in weanling male rats which became relatively shorter in length and narrower in girth as they grew, compared to controls⁷. In mouse embryos and fetuses maternal exposure to toluene during pregnancy was associated with a higher incidence of the presence of 14 ribs, suggesting the teratogenic toxicity of toluene on the skeletal system⁸. In two other studies, delayed ossification of sternbrae after inhalation of benzene in rats⁹ and skeletal variations due to oral or inhalational benzene exposure in mice and rabbits¹⁰ were reported. In a study conducted at a forensic medical institute, where bone age measurement is routinely performed in determination of the identities of prosecuted glue vapor abusers, a significant bone age retardation has been demonstrated in this population when compared to healthy controls¹¹. The present study is the first one

Table I. Comparison of the Data Obtained in the Study and Control Groups*

	Study group (n=25)	Control group (n=30)	p value
Chronological age (years)	14.9 ± 2.5	15.1 ± 2.9	$p > 0.05$
Height (cm)	162.3 ± 11.2	165.1 ± 10.8	$p > 0.05$
Weight (kg)	53.3 ± 8.9	54.7 ± 9.4	$p > 0.05$
Serum calcium (mg/dl)	9.41 ± 0.7	9.35 ± 0.5	$p > 0.05$
Serum phosphorus (mg/dl)	5.23 ± 0.5	5.19 ± 0.9	$p > 0.05$
Alkaline phosphatase (mU/ml)	251 ± 25	249 ± 21	$p > 0.05$
Bone mineral density (g/cm^2)	0.7065 ± 0.116	0.8755 ± 0.182	$p = 0.001$

* Values are given as mean \pm SD.

investigating the bone mineral content by DXA method and reporting a significantly reduced BMD in adolescents who chronically abused glue vapor. DXA is a new method that permits BMD to be measured accurately and directly. Lumbar spinal BMD measurements are preferred in this method. The spine is formed mainly of trabecular bone, which has higher surface-to-volume ratio and is more active metabolically than other regions of the skeleton. Because of the greater rapidity of bone turnover in the trabecular compartment than in the cortical one, the spine is thought to be a more sensitive site than the long bones of the upper and lower extremities for evaluating the effects of various stimuli on bone mineral status of the body^{5,6,12}.

Bone mineral content is influenced by genetic, hormonal and exogenous factors such as physical activity, diet, certain medications, and exposure to sunlight. Cases in both the study and control groups in our study had relatively the same degree of physical activity and were living in the same region, being exposed to relatively the same amount of sunlight, although we could not determine the objective degrees and/or amounts of these two parameters. Moreover, anthropometric measurements and serum calcium, phosphorus and ALP, which indirectly reflect growth, development and nutritional status, were not significantly different between the study and control groups in this study.

The other factors which might possibly affect bone mineralization in this study were alcohol consumption and smoking. However, there are some controversies about the effects of these habits on bone mineral density values. A detrimental effect of chronic alcohol abuse on the skeleton has been shown in two studies^{13,14}, whereas two other studies^{15,16} have suggested a beneficial effect of light-to-moderate alcohol consumption, and some other studies¹⁷⁻¹⁹ have demonstrated no significant effects. Regarding smoking, some negative effects on the skeletal system have been reported^{15,17}, whereas two other studies^{18,19} found no obvious effects on bone mineralization. Most of the cases in our study group were smokers and consumed alcohol although we could not quantify amounts. Thus, to demonstrate the exclusive effects of glue sniffing on bone mineralization, it would certainly be more appropriate, if

possible, to compare our study group with a control group consisting of cases who merely smoked and consumed alcohol but did not abuse any volatile substances.

Results of the present study indicate that teenagers with glue vapor abuse may be at risk for developing osteopenia, possibly due to the toxic effects of glue vapor on the skeletal system. BMD peaks by the age of 20, makes a long plateau, and decreases after the age of 40. A lower peak bone mass in adolescence has been suggested to be associated with a greater risk of osteoporosis and fracture in the future^{20,21}. Teenagers with glue vapor abuse, therefore, may carry an increased risk of future fracture even though the exact mechanism(s) responsible for the toxicity of glue vapor on bone metabolism remains to be determined. Commercially available glue products generally contain toluene, benzene, xylene, trichloroethylene, tetrachloroethylene, methylene chloride, trichlorethane, carbon tetrachloride, acetone, naphtha, and n-hexane. To determine the exact component of glue responsible for bone demineralisation may be of value in proposing a change in the composition of the glue. Education and/or rehabilitation programs currently have the greatest importance in preventing and overcoming the harmful effects of this public health problem which is so common in young children and adolescents.

REFERENCES

1. Inhalant abuse. American Academy of Pediatrics, Committee on Substance Abuse and Committee on Native American Child Health. *Pediatrics* 1996; 97: 420-423.
2. Hyaden JW, Peterson RG, Bruckner JV. Toxicology of toluene (methylbenzene). Review of current literature. *Clin Toxicol* 1977; 11: 549-559.
3. Musclow CE, Awen CF. Glue sniffing: report of a fatal case. *Can Med Assoc J* 1971; 104: 315-319.
4. Streicher HZ, Gabow PA, Moss AH, Kono D, Kaehny WD. Syndromes of toluene sniffing in adults. *Ann Intern Med* 1981; 94: 758-762.
5. Kroger H, Kotaniemi A, Vainio P, Alhava E. Densitometry of the spine and femur in children by dual-energy X-ray absorptiometry. *Bone Miner* 1992; 17: 75-85.
6. Akın R, Okutan V, Sarıcı Ü, Altunbaş A, Gökçay E. Evaluation of bone mineral density in children receiving antiepileptic drugs. *Pediatr Neurol* 1998; 19: 129-131.
7. Pryor GT. A toluene-induced motor syndrome in rats resembling that seen in some human solvent abusers. *Neurotoxicol Teratol* 1991; 13: 387-400.
8. Shigeta S, Aikawa H, Misawa T. Effects of maternal exposure to toluene during pregnancy on mouse embryos and fetuses. *Tokai J Exp Clin Med* 1982; 7: 265-270.

9. Green JD, Leong BK, Laskin S. Inhaled benzene fetotoxicity in rats. *Toxicol Appl Pharmacol* 1978; 46: 9-18.
10. Murray FJ, John JA, Rampy LW, Kuna RA, Schwetz BA. Embryotoxicity of inhaled benzene in mice and rabbits. *Am Ind Hyg Assoc J* 1979; 40: 993-998.
11. Yorulmaz C, Dünderöz R, Arıcan N, Denli M, Tuğcu H. Uçucu madde kötüye kullanımında görülen organ hasarı ve kemik yaşı değişiklikleri. *Adli Tıp Dergisi* 2001; 15: 6-13.
12. Southard RN, Morris JD, Mahan JD, et al. Bone mass in healthy children: measurement with quantitative DXA. *Radiology* 1991; 179: 735-738.
13. Wezeman FH, Emanuele MA, Emanuele NV, et al. Chronic alcohol consumption during male rat adolescence impairs skeletal development through effect on osteoblast gene expression, bone mineral density, and bone strength. *Alcohol Clin Exp Res* 1999; 23: 1534-1542.
14. Sampson HW, Perks N, Champney TH, DeFee B 2nd. Alcohol consumption inhibits bone growth and development in young actively growing rats. *Alcohol Clin Exp Res* 1996; 20: 1375-1384.
15. Baron JA, Farahmand BY, Weiderpass E, et al. Cigarette smoking, alcohol consumption, and risk of hip fracture in women. *Arch Intern Med* 2001, 161: 983-988.
16. Sampson HW, Gllager S, Lange J, Chondra W, Hogan HA. Binge drinking and bone metabolism in a young actively growing rat model. *Alcohol Clin Exp Res* 1999, 23: 1228-1231.
17. Burger H, deLaet CE, van Daele PL, et al. Risk factors for increased bone loss in an elderly population: the Rotterdam study. *Am J Epidemiol* 1998; 147: 871-879.
18. Orozco P, Nolla JM. Associations between body morphology and bone mineral density in premenopausal women. *Eur J Epidemiol* 1997; 13: 919-924.
19. Kroger H, Laitinen K. Bone mineral density measured by dual-energy X-ray absorptiometry in normal men. *Eur J Clin Invest* 1992; 22: 454-460.
20. Matkovic V, Jelic T, Wardlaw GM, et al. Timing of peak bone mass in Caucasian females and its implication for the prevention of osteoporosis. Inference from a cross-sectional model. *J Clin Invest* 1994; 93: 799-808.
21. Melton LJ 3rd, Kan SH, Wahner HW, Riggs BL. Lifetime fracture risk: an approach to hip fracture risk assessment based on bone mineral density and age. *J Clin Epidemiol* 1988; 41: 985-994.